


The Open COVID Journal

Content list available at: <https://opencovidjournal.com>



RESEARCH ARTICLE

The Impact of COVID-19 Strict Lockdown on the Air Quality of Smart Cities of Rajasthan, India

Tansukh Barupal¹, Praveen Kumar Tak², Mukesh Meena^{3,*} , Pradeep Kumar Vishwakarma⁴ and Prashant Swapnil⁵

¹Sangidas Balkishan Kothari Government PG College, Jaisalmer – 345001, Rajasthan, India

²M.L.V. Government College, Bhilwara – 311001, Rajasthan, India

³Department of Botany, Laboratory of Phytopathology and Microbial Biotechnology, Mohanlal Sukhadia University, Udaipur – 313001, Rajasthan, India

⁴Department of Mathematics and Statistics, Mohanlal Sukhadia University, Udaipur – 313001, Rajasthan, India

⁵Department of Botany, School of Biological Science, Central University of Punjab, Bhatinda, Punjab – 151401, India

Abstract:

Aim:

The main focus of this study is to evaluate the air quality by comparing the concentration of particulate matter PM_{2.5}, PM₁₀, NO₂, CO, SO₂ and ozone of smart cities of Rajasthan before the lockdown and during the period of lockdown.

Background:

In India, the first case of the COVID-19 was reported on January 30th, 2020. Indian government declared strict lockdown, *i.e.*, public health emergency in India on March 24th, 2020, which is implemented from March 25th, 2020, to April 14th, 2020, for 21 days.

Objective:

The objective of this study is to evaluate the air quality by comparing the levels of all parameters of air pollution during the COVID-19 lockdown period with values registered in the pre-lockdown period.

Methods:

Data were obtained from four automatic monitoring stations under the control of the Central Pollution Control Board, New Delhi (<https://www.cpcb.nic.in/>). Data regarding all the parameters were recorded as 24 hours average period.

Results:

CO levels showed the highest significant reduction in Udaipur (50.76%) followed by Jaipur (19.96%), Ajmer (17.11%), and Kota (5.51%) due to the ban on transport and driving. The levels of PM_{2.5}, PM₁₀, NO₂, and SO₂ were also decreased substantially for each smart city. Ozone concentrations were recorded greater than before due to decreased nitrogen oxides levels.

Conclusion:

This study can be useful considering our present role in environmental restoration or environmental destruction. It will also be helpful in updating our present plan toward the assurance and conservation of nature.

Keywords: COVID-19, Sulfur dioxide, PM, Ozone, Public health, Lockdown.

Article History

Received: September 08, 2021

Revised: December 9, 2021

Accepted: January 6, 2022

1. INTRODUCTION

On December 8th, 2019, the novel SARS-CoV-2 coronavirus first reported in Wuhan, China. It causes COVID-19 dis-

ease, which spreads across the world. On January 9th, 2020, coronavirus was discovered as a novel infectious agent of COVID-19 [1, 2]. It is a communicable disease that spreads very quickly from one person to another. To date, all over the world, the total reported cases are estimated to be 4,628,821, and 308,654 people died because of this pandemic. All world governmental bodies are worried about this crisis because its cure has not been discovered yet. Currently, prevention is the

* Address correspondence to this author at Department of Botany, Laboratory of Phytopathology and Microbial Biotechnology, Mohanlal Sukhadia University, Udaipur – 313001, Rajasthan, India;
E-mails: mukeshmeenamsu@gmail.com, mukeshmeenabhu@gmail.com

only way to avoid infection. For this, many countries announced the complete lockdown to prevent the spread of infection and succeeded. The Indian government also imposed total five lockdown with various rules and regulations. Out of five lockdowns, the first lockdown implemented from March 25th, 2020, to April 14th, 2020, for 21 days was very strict; there were strict restrictions on most of the activities, like the transportation, market, industries, etc.

Rajasthan also implemented all rules and regulations of this lockdown. We know that industries and transportation are major sources of pollutants, like particulate matter (PM_{2.5}), nitrogen dioxide (NO₂), sulfur dioxide (SO₂), carbon monoxide (CO), and ozone (O₃), which are directly impacting the air quality. It is a major concern of health risk and cause of large premature mortality [3, 4]. In research, it has been found that the air quality parameters of the most polluted cities (Delhi, Lucknow, Kolkata, Patna, and Bihar) and the least polluted cities (Amaravati, Panchkula, Shillong, and Chennai) of India have been changed during the lockdown period. The information regarding air quality parameters of the most and least polluted cities is provided in Table 1 [5]. In India, Rajasthan is also a very growing state with a lot of industrial development, power plants, and transportation. According to the WHO Global Air Pollution Database 2018, all four smart cities, *i.e.*, Jaipur, Udaipur, Ajmer, and Kota, are listed in the polluted cities of the world. The major pollutants include CO, which is mainly introduced into the air by the burning of fossil fuels and vehicles exhaust, SO₂ from petroleum refineries and thermal power plants, NO₂ from thermal power plants and vehicle exhausts, ozone from industrial release and vehicle exhaust, and particulate matter consists of solid in the air in the

form of dust, smoke, and vapour.

Recently, researchers have reported that the lockdown period dealt with social distancing and decreasing vehicle transport, significantly improving air quality. Tobias *et al.* [6] reported that NO₂ and particulate matter (PM₁₀) were reduced by 51% and 31%, respectively, in Barcelona (Spain) during the lockdown. Nakada and Urban [7] analyzed that concentrations of CO, NO, and NO₂ decreased by 64.8%, 77.3%, and 54.3%, respectively, during the partial lockdown in the Sao Paulo state of Brazil. Dantas *et al.* [8] also reported a significant reduction in NO₂, CO, and PM₁₀ levels during the COVID-19 partial lockdown in Rio de Janeiro, Brazil. It has been estimated that long-term or wide exposure to air pollution is correlated with respiratory diseases, such as bronchitis, aggravated asthma, respiratory allergies, heart disease, stroke, and death [9 - 11]. Particulate matters (PM_{2.5} and PM₁₀) are considered as one of the most health risk factors and cause millions (~9%) of deaths around the world annually [11]. A higher concentration of NO₂ emitted from transportation and fuel combustion (outdoor anthropogenic sources) is another most important air pollutant that causes toxicity to human respiratory systems [12, 13]. A higher concentration of NO₂ causes severe illnesses, like cytokine-mediated lungs inflammation, diabetes, hypertension, cardiovascular diseases, and even death [11, 14]. Polluted air is associated with more SARS fatality and influenza [15, 16]. *In situ* study declared that SARS-CoV-2 shows stability in ambient aerosols, which can be considered a source of COVID-19 transmission [17]. Atmospheric aerosol works as a cofactor that can alter immune defence response system pathways and causes severe pneumonia [16, 18 - 20].

Table 1. Comparison of average air quality parameters before and during lockdown phase among (A) Major polluted cities and (B) Least polluted cities of India [5].

(A)								
Air quality Parameters	Delhi, Delhi		Lucknow, Uttar Pradesh		Kolkata, West Bengal		Patna, Bihar	
	Before Lockdown	During Lockdown	Before Lockdown	During Lockdown	Before Lockdown	During Lockdown	Before Lockdown	During Lockdown
PM _{2.5}	151.8	54.8	245	186.66	137.7	63.9	117.5	29.4
PM ₁₀	166.0	71.9	NA	NA	109.5	72.0	226	NA
NO ₂	60.6	21.9	37	19.33	28.1	9.6	67.5	60
CO	91.4	61.3	61.66	46.66	23.5	16.8	45	29
SO ₂	13.4	12.0	5.66	5.66	10.7	8.2	9	7
Ozone	30.1	45.4	35	31.66	40.2	42.8	71	70

(B)								
Air quality Parameters	Amaravati, Andhra Pradesh		Panchkula, Haryana		Shillong, Meghalaya		Chennai, Tamil Nadu	
	Before Lockdown	During Lockdown	Before Lockdown	During Lockdown	Before Lockdown	During Lockdown	Before Lockdown	During Lockdown
PM _{2.5}	40.5	29	32	23.33	59	74.5	38.5	25.33
PM ₁₀	42	36.66	NA	NA	50	66	NA	NA
NO ₂	11.5	5.3	21.5	22.66	3.5	3	4	2
CO	25.5	10.66	2	21.33	16	14.5	16	38.66
SO ₂	26	23	14	9.33	6	9	7.5	8.66
Ozone	29	20.66	56	35	52.5	11.5	52.5	24.33

In India, indoor air pollution is also causing serious public health issues because most households depend on the use of solid fuel. The long periods of lockdown to control COVID-19 in India causes adverse health effects through household air pollution [21]. In this study, we have selected four major cities of Rajasthan, *i.e.*, Jaipur, Ajmer, Udaipur, and Kota. The details of the selected cities are provided in Table 2 [22]. The main goal of this study is to analyze the impact of strict lockdown amid COVID19 on the air quality of the smart cities of Rajasthan, India, by comparing the particulate matter, carbon monoxide, nitrogen dioxide, sulfur dioxide, and ozone concentrations determined during the strict lockdown with values obtained before and during the lockdown.

2. MATERIAL AND METHODS

2.1. Studied Area

Rajasthan is the largest state by area in India. It covers an area of 342,239 square kilometers (132,139 sq mi) or 10.4% of the total geographical area of India. It is located on the western side of the country, situated between 23 30' and 30 11' North latitude and 69 29' and 78 17' East longitude. In this study, we have selected four smart cities, *i.e.*, Jaipur, Udaipur, Ajmer, and Kota, as shown in Fig. (1), and their geographical locations, coordinates, and monitoring station are shown in Table 2 [22].

Jaipur, a smart city, is situated in the East-central part of

the state. It is the capital and largest city of Rajasthan. Udaipur, Ajmer, Kota are also important cities of Rajasthan. Udaipur is located in the southern region of Rajasthan. Ajmer is situated in the center or heart of Rajasthan, and the Kota is situated at the center of the southeastern region of Rajasthan. The main characteristics are briefly described in Table 2 [22].

2.2. Experimental Data and Analysis

In this study, we have selected a total of four parameters, *i.e.*, $PM_{2.5}$, NO_2 , SO_2 , CO , and ozone concentration, to assess the air quality of smart cities of Rajasthan. These data were obtained from four automatic monitoring stations, *i.e.*, Adarsh Nagar, Jaipur – RSPCB, Ashok Nagar, Udaipur – RSPCB, Civil Lines, Ajmer – RSPCB and Shrinath Puram, Kota – RSPCB under the control of Central Pollution Control Board, New Delhi (<https://www.cpcb.nic.in/>). Data of all parameters were recorded as 24 hours average period, and the air quality of these cities was compared with the standard air quality index (AQI) before and during the lockdown (Table 3) [5]. Experimental data obtained by each monitoring station were analyzed using standard statistical techniques. Comparisons of each parameter, *i.e.*, obtained during the lockdown and before the lockdown, were made by graphical representation. This study also included the data regarding total COVID-19 cases and death (mortality) registered in India in pre- and during the lockdown period (Table 4) [23].




Fig. (1). Geographical status of the studied smart cities of Rajasthan situated in India.

Table 2. Description of the studied smart cities of Rajasthan, India [22].

S. No.	Smart city	Coordinates	Area (km ²)	City Population	The Monitoring Station
1.	Jaipur	26.9124°N, 75.7873° E	484.6	3,046,163	Adarsh Nagar, Jaipur - RSPCB
2.	Udaipur	24.5854°N, 73.7125° E	37	451,100	Ashok Nagar, Udaipur - RSPCB
3.	Ajmer	26.4499°N, 74.6399° E	55.76	542,321	Civil Lines, Ajmer - RSPCB
4.	Kota	25.2138°N, 75.8648° E	527	1,001,694	Shrinath Puram, Kota - RSPCB

Table 3. (A) Standard AQI and (B) AQI registered in smart cities of Rajasthan in pre and during lockdown period [5].

(A)		
AQI	Remark (Air Pollution Level)	Possible Health Impacts
0-50	Good	Minimal impact
51-100	Satisfactory	Minor breathing discomfort to sensitive people
101-200	Moderate	Breathing discomfort to the people with lungs, asthma, and heart diseases
201-300	Poor	Breathing discomfort to most people on prolonged exposure
301-400	Very Poor	Respiratory illness on prolonged exposure
401-500	Severe	Affects healthy people and seriously impacts those with existing diseases

(B)			
S. No.	Smart City	Air Quality Index (AQI)	
		Pre-Lockdown Period	During Lockdown Period
1.	Jaipur	91.00	66.33
2.	Udaipur	81.00	53.33
3.	Ajmer	102.50	58.00
4.	Kota	95.00	73.67

Table 4. COVID-19 cases and mortality registered in India in pre- and during lockdown period [23].


Duration	Date	Cases	Mortality	Percent (%)	Average Percent (%)
Pre-Lockdown Period	March 1	3	0	0	1.81
	March 5	30	0	0	
	March 10	62	0	0	
	March 15	114	2	1.75	
	March 20	249	5	2.00	
	March 24	536	11		
During Lockdown Period	March 25	657	13	1.97	2.70
	March 30	1251	33	2.63	
	April 1	4289	59	1.37	
	April 5	4778	119	2.49	
	April 10	7600	250	3.28	
	April 14	11887	349	2.94	

3. RESULTS AND DISCUSSION

Changes in concentrations of major pollutants during the pre-lockdown and during-lockdown period are depicted in Fig. (2). Indian government declared a strict lockdown in India on March 24th, 2020, which was implemented from March 25th, 2020, to April 14th, 2020. During this period, *i.e.*, 21 days, all four smart cities of Rajasthan observed substantial diminution of the pollutants. The present article assessed air quality during the strict lockdown (March 25th, 2020 to April 14th, 2020).

Results showed the highest reduction in PM_{2.5} found in Ajmer (53.07%), followed by Jaipur (38.32%), Kota (28.39%), and Udaipur (15.37%), respectively. PM₁₀ was also found to be reduced in Ajmer (45.56%), followed by Jaipur (36.18%), Udaipur (29.40%), and Kota (19.69%), respectively. Whereas, concerning NO₂, the highest reduction was found in Udaipur (66.07%), followed by Jaipur (62.90%), Kota (56.83%), and Ajmer (54.36%), respectively. In the case of SO₂, the highest reduction was found in Udaipur (28.70%), and the lowest was found in Jaipur (11.71%).

(A). Before lockdown



(B). During lockdown




Fig. (2 A,B). Comparative analysis of different air quality parameters (PM_{2.5}, PM₁₀, NO₂, CO, SO₂ and ozone) of smart cities of Rajasthan, i.e., Jaipur, Udaipur, Ajmer, and Kota before and during the period of lockdown.

However, increasing O₃ concentration was recorded at 12.82% in Kota and 6.08% in Ajmer, but Udaipur and Jaipur showed negative results -6.08 and -13.59%, respectively. Overall the result concluded that the air quality index of the cities was found more satisfactory during the lockdown period than the pre-lockdown period by comparing with standard AQI. In India, a high number of people were infected due to coronavirus. During the pre-lockdown (from March 1st to 24th, 2020), a total of 994 active cases and 18 deaths were registered. During the lockdown period (March 25th, 2020, to April 14th, 2020), a total of 30462 active cases and 823 deaths were registered. It was reported that active cases were increased during lockdown conditions (Table 4). The average death rate percentage was found to be 1.81% during the pre-lockdown and 2.70% during the lockdown phase. The relationship between active cases and deaths may specify the severity of any pandemic.

It has been investigated that meteorological conditions, such as air, temperature, wind direction, and wind speed, are known to greatly impact urban air pollutant’s level *via* some physical and chemical processes, like accumulation or dispersion and multiphase reactions of aerosol formation and growth [24]. Therefore, it is reasonable to suppose that changes in urban air pollutant concentrations during the COVID-19 lockdown may partially depend on meteorological conditions during that period [25]. Movements of air affect the fate of air pollutants. Concentrations of pollutants build up by non-dispersion of air pollutants due to calm air. However, when strong, turbulent winds blow, the concentration of air pollutants decreases. To estimate air quality, some important factors need to be measured, which can help in understanding the chemical reactions which occur in atmospheres, such as wind speed and direction, temperature, humidity, rainfall, and solar radiation [26].

In India, Rajasthan is also a growing state with the establishment of various kinds of industries and high pressure of transportation. Moreover, WHO also listed all four smart cities, *i.e.*, Jaipur, Udaipur, Ajmer, and Kota, as the most polluted cities of the world based on the global air pollution database. The air quality of all these smart cities was measured by the data provided by the four automatic monitoring stations. The impact of lockdown on air quality in different parts of the world indicated how human beings drastically affect the environment. Therefore, the results of the present article will possibly help us in rethinking the kind of future we are establishing. The article may also help explain how industries and transportation negatively affect the environment and how we can restore the environment and provide a good quality ecosystem to future generations. Currently, on the one hand, the outrageous viruses have affected our lives, and on the other hand, the process of environmental restoration is also going on. Hence, global concern for air pollution has led to drawing significant attention towards analyzing air pollution during the pandemic. In the metropolitan area of São Paulo (2100 km³), the NO₂ concentration reduction is visualized by satellite measurements [5, 27, 28]. However, Sharma *et al.* [29] reported an increase of 17% in O₃ concentration in India. In China, about 25% carbon and 30% NO₂ emission have been reduced during the period of lockdown [30 - 33]. Overall, pollutant concentrations decreased in all analyzed areas.

CONCLUSION

This study can be helpful in evaluating our present role in environmental restoration or environmental destruction. It will also be helpful in updating the present plan toward the assurance and conservation of nature. The main focus of this study is to evaluate the air quality by comparing the concentration of particulate matter PM_{2.5}, PM₁₀, NO₂, CO, SO₂, and ozone in smart cities of Rajasthan, *i.e.*, Jaipur, Udaipur, Ajmer, and Kota before and during the period of lockdown. CO levels showed the highest significant reduction in Udaipur (50.76%), followed by Jaipur (19.96%), Ajmer (17.11%), and Kota (5.51%), due to the ban on transport and driving. The levels of PM_{2.5}, PM₁₀, NO₂, and SO₂ were also found to be substantially decreased for each smart city.

Ozone concentration was recorded greater than before lockdown due to decreased nitrogen oxides level. Although, more specific studies are required to explore the interactions between SARS-CoV-2 and air pollutants to understand their impacts on human health. Future studies should be investigated to examine the role of indoor air pollution (biomass and tobacco smoke) during COVID-19. Biology and atmospheric chemistry should be implicated in a more holistic way to the disease management and mitigation to address the current pandemic condition and future viral epidemics. We should not ignore our surrounding environment, which is causing not only chronic diseases but also severe infectious diseases. To establish a healthy future, we need to reduce air pollution from all sources, such as road traffic and heat generation, by strengthening public health policies.

LIST OF ABBREVIATIONS

PM	=	Particulate Matter
NO ₂	=	Nitrogen Dioxide
SO ₂	=	Sulfur Dioxide
CO	=	Carbon Monoxide
NO	=	Nitric Oxide
WHO	=	World Health Organization
AQI	=	Air Quality Index

AUTHORS' CONTRIBUTION

TSB, MM, and PKT contributed to the study concept and design. PKT, TSB, and MM participated in data collection. Data analysis was performed by PKV and MM. TSB, PS, and MM contributed to the writing of the manuscript.

ETHICS APPROVAL AND CONSENT TO PARTICIPATE

Not applicable.

HUMAN AND ANIMAL RIGHTS

No animals/humans were used in the studies that are the basis of this research.

CONSENT FOR PUBLICATION

Not applicable.

AVAILABILITY OF DATA AND MATERIALS

Not applicable.

FUNDING

None.

CONFLICT OF INTEREST

The authors declare no conflict of interest, financial or otherwise.

ACKNOWLEDGEMENTS

The authors would like to thank the Central Pollution Control Board, Ministry of Environment, Forest and Climate Change (MoEFCC), and Ministry of Human Resources and Development, Government of India.

REFERENCES

- [1] Cheng VCC, Wong SC, To KKW, Ho PL, Yuen KY. Preparedness and proactive infection control measures against the emerging novel coronavirus in China. *J Hosp Infect* 2020; 104(3): 254-5. [<http://dx.doi.org/10.1016/j.jhin.2020.01.010>] [PMID: 31962139]
- [2] Swapnil P, Meena M. SARS-CoV-2 (COVID-19) pandemic outbreak: A worldwide challenge and contemporary situation. *Virology* 2021; 10: 201.
- [3] Sharma S, Zhang M, Anshika, Gao J, Zhang H, Kota SH. Effect of restricted emissions during COVID-19 on air quality in India. *Sci Total Environ* 2020; 728: 138878. [<http://dx.doi.org/10.1016/j.scitotenv.2020.138878>] [PMID: 32335409]
- [4] Swapnil P, Meena M, Barupal T, Sompura Y, Hada D. Structure, transmission, diagnostic symptoms, host and entry mechanism of COVID-19: A review. *Coronaviruses* 2021; 2(5): 1-9.

- [5] National Air Quality Index - CPCB. Available from: https://app.cpcbcr.com/AQI_India/
- [6] Tobias A, Carnero C, Reche C, *et al.* Changes in air quality during the lockdown in Barcelona (Spain) one month into the SARS-CoV-2 epidemic. *Sci Total Environ* 2020; 726: 138540. [<http://dx.doi.org/10.1016/j.scitotenv.2020.138540>] [PMID: 32302810]
- [7] Nakada LYK, Urban RC. COVID-19 pandemic: Impacts on the air quality during the partial lockdown in São Paulo state, Brazil. *Sci Total Environ* 2020; 730: 139087. [<http://dx.doi.org/10.1016/j.scitotenv.2020.139087>] [PMID: 32380370]
- [8] Dantas G, Siciliano B, França BB, da Silva CM, Arbilla G. The impact of COVID-19 partial lockdown on the air quality of the city of Rio de Janeiro, Brazil. *Sci Total Environ* 2020; 729: 139085. [<http://dx.doi.org/10.1016/j.scitotenv.2020.139085>] [PMID: 32361428]
- [9] WHO. World Health Organization. World Health Organization Emergency Dashboard 2020. Available from: <https://covid19.who.int/region/searo/country/in>
- [10] MHA (Ministry of Home Affairs). Govt of India Circulars for Covid-19 2020. Available from: <https://www.mha.gov.in/notifications/circulars-covid-19>
- [11] Hu B, Guo H, Zhou P, Shi ZL. Characteristics of SARS-CoV-2 and COVID-19. *Nat Rev Microbiol* 2021; 19(3): 141-54. [<http://dx.doi.org/10.1038/s41579-020-00459-7>] [PMID: 33024307]
- [12] Chen LA, Chien LC, Li Y, Lin G. Nonuniform impacts of COVID-19 lockdown on air quality over the United States. *Sci Total Environ* 2020; 745: 141105. [<http://dx.doi.org/10.1016/j.scitotenv.2020.141105>] [PMID: 32731074]
- [13] Berman JD, Ebisu K. Changes in U.S. air pollution during the COVID-19 pandemic. *Sci Total Environ* 2020; 739: 139864. [<http://dx.doi.org/10.1016/j.scitotenv.2020.139864>] [PMID: 32512381]
- [14] Unnithan PSG. Kerala reports first Coronavirus case in India 2020. Available from: <https://www.indiatoday.in/india/story/kerala-reports-first-confirmed-novel-coronavirus-case-in-india-1641593-2020-01-30>
- [15] He G, Pan Y, Tanaka T. COVID-19, city lockdowns, and air pollution: evidence from China. *medRxiv* 2020. [<http://dx.doi.org/10.1101/2020.03.29.20046649>]
- [16] Ndegwa S. An environmental silver lining amid COVID-19 cloud. China global television network, opinion. 2020. Available from: <https://news.cgtn.com/news/2020-06-06/An-environmental-silver-linings-amid-COVID-19-cloud-R4TpEvr0ty/index.html>
- [17] Zambrano-Monserrate MA, Ruano MA, Sanchez-Alcalde L. Indirect effects of COVID-19 on the environment. *Sci Total Environ* 2020; 728: 138813. [<http://dx.doi.org/10.1016/j.scitotenv.2020.138813>] [PMID: 32334159]
- [18] Meena M, Swapnil P. A review of contagious coronavirus (SARS-Cov-2) their clinical features, diagnosis, preventions and treatment. *J Hum Virol Retrovirol* 2020; 8(4): 99-105. [<http://dx.doi.org/10.15406/jhvr.2020.08.00227>]
- [19] European Environmental Agency. Air pollution goes down as Europe takes hard measures to combat 2020. Available from: <https://www.eea.europa.eu/highlights/air-pollution-goes-down-as>
- [20] Garg A, Kumar A, Gupta NC. Impact of lockdown on ambient air quality in COVID-19 affected hotspot cities of India: Need to readdress air pollution mitigation policies. *Environ Claims J* 2021; 1(1): 65-76. [<http://dx.doi.org/10.1080/10406026.2020.1822615>]
- [21] Rani R, Arokiasamy P, Sikarwar A. Household air pollution during COVID-19 pandemic: A concern in India. *J Public Aff* 2021; 21(4): e2729. [<http://dx.doi.org/10.1002/pa.2729>] [PMID: 34512188]
- [22] Census 2011 India. Available from: <https://www.indiatoday.in/india/story/kerala-reports-first-confirmed-novel-coronavirus-case-in-india-1641593-2020-01-30>
- [23] Worldometers. Available from: <https://www.worldometers.info/coronavirus/country/india/>
- [24] Cai WJ, Wang HW, Wu CL, Lu KF, Peng ZR, He HD. Characterizing the interruption-recovery patterns of urban air pollution under the COVID-19 lockdown in China. *Build Environ* 2021; 205: 108231. [<http://dx.doi.org/10.1016/j.buildenv.2021.108231>] [PMID: 34393324]
- [25] Diaz-Avalos C, Juan P, Chaudhuri S, Sáez M, Serra L. Association between the new COVID-19 cases and air pollution with meteorological elements in nine counties of New York state. *Int J Environ Res Public Health* 2020; 17(23): 9055. [<http://dx.doi.org/10.3390/ijerph17239055>] [PMID: 33291673]
- [26] Queensland Government. Available from: <https://www.qld.gov.au>
- [27] Gorelick N, Hancher M, Dixon M, Ilyushchenko S, Thau D, Moore R. Google earth engine: planetary-scale geospatial analysis for everyone. *Remote Sens Environ* 2017; 202: 18-27. [<http://dx.doi.org/10.1016/j.rse.2017.06.031>]
- [28] Yuen KY, Chan PK, Peiris M, *et al.* Clinical features and rapid viral diagnosis of human disease associated with avian influenza A H5N1 virus. *Lancet* 1998; 351: 467e71. [[http://dx.doi.org/10.1016/S0140-6736\(98\)01182-9](http://dx.doi.org/10.1016/S0140-6736(98)01182-9)]
- [29] Sharma S, Swapnil P, Meena M. A novel 2019 severe acute respiratory syndrome coronavirus 2 (COVID-19) as global pandemic. *Cellular Life Sci J* 2020; 5(2): 000154. [<http://dx.doi.org/10.23880/cclsj-16000154>]
- [30] He G, Pan Y, Tanaka T. The short-term impacts of COVID-19 lockdown on urban air pollution in China. *Nat Sustain* 2020; 3(12): 1005-11. [<https://doi.org/10.1038/s41893-020-0581-y>]. [<http://dx.doi.org/10.1038/s41893-020-0581-y>]
- [31] Liu F, Page A, Strode SA, *et al.* van der AR, Veeffkind P, Levelt PF, Hauser OP, Joiner J. Abrupt declines in tropospheric nitrogen dioxide over China after the outbreak of COVID-19. *Sci Adv* 2020; 6(28): eabc2992. [<http://dx.doi.org/10.1126/sciadv.abc2992>] [PMID: 32923601]
- [32] Barupal T, Tak PK, Meena M. COVID-19: Morphology, characteristics, symptoms, prevention, clinical diagnosis and current scenario. *Coronaviruses* 2020; 1(1): 82-9. [<http://dx.doi.org/10.2174/2666796701999200617161348>]
- [33] Cheng VC, Lau SK, Woo PC, Yuen KY. Severe acute respiratory syndrome coronavirus as an agent of emerging and reemerging infection. *Clin Microbiol Rev* 2007; 20: 660e94. [<http://dx.doi.org/10.1128/CMR.00023-07>]